

BRESKVA I NEKTARINA

ORIJENTACIJSKI PROGRAM ZAŠTITE

VRIJEME PRIMJENE	PRIPRAVAK	SUZBIJA
POČETAK OPADANJA LIŠĆA	■ CUPRABLAU Z 0,8% (800 g na 100 l vode)	kovrčavost i rupičavost lista
	■ FERRUS 648 30-80 g/biljci	Prevenција i uklanjanje simptoma nedostatka željeza (ferokloroza)
 OD MIROVANJA DO VIDLJIVIH ČAŠKI	■ CUPRABLAU Z 0,5% (500 g na 100 l vode)	kovrčavost i rupičavost lista
	+ ■ mineralno ulje	zimске stadije štetnika
	■ HERKULES SUPER 4 - 8 l/ha ili ■ OXALIS 3 - 6 l/ha	jednogodišnje i višegodišnje širokolisne i uskolisne korove
 POJAVA CVJETOVA	■ CHROMODIN S-65 0,1 - 0,12% (100 - 120 g na 100 l vode) ili ■ DELAN 700 WDG 0,075 - 0,1% (75 - 100 g na 100 l vode) ili ■ ZIRAM 76 WG 2,5 kg/ha (250 g na 100 l vode)	kovrčavost lista breskve
	+ ■ TEPPEKI 500 WG 0,12 - 0,14 kg/ha (12 - 14 g na 100 l vode)	zelenу breskvinu uš
 POČETAK CVATNJE	■ SIGNUM 0,056 - 0,075% (56 - 75 g na 100 l vode)	moniliju (sušenje rodniх grančica), palež cvijeta
 ZAVRŠETAK CVATNJE	■ SIGNUM 0,056 - 0,075% (56 - 75 g na 100 l vode)	moniliju (sušenje rodniх grančica), palež cvijeta
	■ DELAN 700 WDG 0,075 - 0,1 % (75-100 g na 100 l vode) ili ■ ZIRAM 76 WG 2,5 kg/ha (250 g na 100 l vode)	kovrčavost lista breskve
	+ ■ TEPPEKI 500 WG 0,12 - 0,14 kg/ha (12 - 14 g na 100 l vode) ili ■ BOXER 200 SL 0,025% (25 ml na 100 l vode)	lisne uši
 FORMIRANI PLOD	■ DELAN 700 WDG 0,075 - 0,1 % (75-100 g na 100 l vode)	šupljikavost lista breskve
	+ ■ CHROMOSUL 80 0,3 - 0,5% (300 - 500 g na 100 l vode)	pepelnicu
 RAST PLODA	■ CHROMODIN S-65 0,1 - 0,12% (100 - 120 g na 100 l vode)	kovrčavost lista
	■ BOXER 200 SL 0,025% (25 ml na 100 l vode) ili ■ CHROMGOR 40 0,1 - 0,15% (100 - 150 ml na 100 l vode)	breskinog savijača, breskinog moljca



NAPOMENA:

Sredstvo CHROMGOR 40 se smije primjenjivati u nasadima bresaka i nektarina najkasnije do 01.01.2015.
Sredstvo Boxer 200 SL tretirati isključivo nakon cvatnje.

■ Herbicidi ■ Fungicidi ■ Insekticidi



OVDJE ČUVAM JA